

Pointwise Bounded Approximation by Polynomials

Anthony G. O'Farrell and Fernando Perez-Gonzalez

Affiliation and addresses:

Anthony G. O'Farrell:

Maynooth College

Co. Kildare

Ireland.

Fernando Perez-Gonzalez:

Dept. de Análisis Matemático

CIMAC

Universidad de la Laguna

38271 La Laguna, Tenerife

Canary Islands

Spain.

Pointwise Bounded Approximation by Polynomials

Anthony G. O'Farrell and Fernando Perez-Gonzalez

1. Introduction.

For a bounded open set $U \subset \mathbf{C}$, we denote by $H^\infty(U)$ the collection of all bounded analytic functions on U . We let X denote $\text{bdy } U$, the boundary of U , Y denote the polynomial hull of U (the complement of the unbounded component of $\mathbf{C} \sim X$), and U^* denote $\text{int } Y$, the interior of Y . We denote the sup norm of a function $f : A \rightarrow \mathbf{C}$ by $\|f\|_A$:

$$\|f\|_A = \sup\{|f(z)| : z \in A\}.$$

We denote the space of all analytic polynomials by $\mathbf{C}[z]$, and we denote the open unit disc by \mathbf{D} and the unit circle by \mathbf{S}^1 .

This paper is about the possibility of approximating a function $f \in H^\infty(U)$, using only polynomials which satisfy the same control as $|f|$ on a prescribed subset $S \subset U$. This kind of approximation was first considered by O.J. Farrell, and has been studied by Rubel and Stray (cf. [4]).

Definition. Let $U \subset \mathbf{C}$ be a bounded open set, and $F \subset \text{clos}(U)$ be closed. We say that (U, F) is a *Farrell pair* if for each $f \in H^\infty(U)$, and each $\epsilon > 0$, there exist $p_m(z) \in \mathbf{C}[z]$, such that

$$\begin{aligned} p_m(z) &\rightarrow f(z), \forall z \in U, \\ \|p_m\|_U &\leq \|f\|_U, \text{ and} \\ \|p_m\|_F &\leq \|f\|_{F \cap U} + \epsilon. \end{aligned}$$

We shall characterise the Farrell pairs.

In view of the Farrell–Rubel–Shields Theorem [2, p.151], $\mathbf{C}[z]$ is pointwise boundedly dense in $H^\infty(U)$ if and only if each $f \in H^\infty(U)$ extends to $H^\infty(U^*)$, and this in

turn happens if and only if there are components U_1, U_2, U_3, \dots of U^* and closed sets E_1, E_2, E_3, \dots such that

$$U = \bigcup_n (U_n \sim E_n)$$

and the sets $E_n \cap U_n$ have inner analytic capacity zero.

Thus, if we are studying situations in which $\mathbf{C}[z]$ is pointwise boundedly dense in $H^\infty(U)$, then there is no loss in generality in assuming that U is a union of some components of U^* (i.e. that the E_n are empty).

We call a set U an *FRS set* if it is open and bounded and is a union of components of its U^* . Note that the components of an FRS set are simply-connected.

Since we lose nothing essential, we shall work exclusively with FRS sets U .

When U is an FRS set, with components U_1, U_2, U_3, \dots , we fix a point $a_n \in U_n$ and let λ_n denote harmonic measure for a_n on $\text{bdy } U_n$. We set $\lambda = \sum_1^\infty \lambda_n / 2^n$, and call λ a *harmonic measure for U* . We note that λ_n is singular to λ_m whenever $n \neq m$ (— the simplest way to see this is to note, first, that U_m and U_n can share at most one common accessible boundary point, second, by Fatou's Theorem, the harmonic measure for U_n is supported on the set of accessible boundary points, and third, harmonic measure contains no point masses). There is a canonical isometric injection $f \mapsto \tilde{f}$ of $H^\infty(U) \rightarrow L^\infty(\lambda)$. There are a number of equivalent ways of describing \tilde{f} . One way is to map the n -th component U_n conformally to \mathbf{D} , transfer $f|_{U_n}$ to a function in $H^\infty(\mathbf{D})$, take nontangential limits a.e. $d\theta$, and transfer them back. This defines \tilde{f} a.e. $d\lambda_n$. Since the various λ_n are mutually-singular, this suffices to define \tilde{f} . A second way is to use Brownian motion. For our present purposes, the most relevant is this. We may choose a sequence p_m of polynomials converging pointwise boundedly on U to f , and converging λ -a.e. See [2, (VI.5.2)]. The limit is \tilde{f} .

For each n , we fix a conformal map $\phi_n : U_n \rightarrow \mathbf{D}$. Even though ϕ_n need not extend continuously to $X_n = \text{bdy}(U_n)$, it does extend as a Borel bijection of a set of full λ_n measure on X_n onto a set of full $d\theta$ measure on \mathbf{S}^1 . The extension is non-unique, but

it is determined up to sets of measure zero at each end. We denote by H_n the image in \mathbf{S}^1 of F under such a bijection.

Theorem 1. *Let U be an FRS set with components U_n ($n = 1, 2, 3, \dots$) and harmonic measure λ . Let F be a closed subset of $\text{clos}(U)$. Then the following are equivalent:*

- (1) (U, F) is a Farrell pair;
- (2) $(U_n, F \cap \text{clos}(U_n))$ is Farrell, for each n .
- (3) For each $f \in H^\infty(U)$,

$$|\tilde{f}| \leq \|f\|_{F \cap U}, \quad \lambda\text{-a.e. on } F;$$

- (4) For each n , $d\theta$ -almost-every point of H_n is a nontangential limit point of $\phi_n(F \cap U_n)$;
- (5) For each closed set $A \subset F \cap \text{bdy}(U)$, there is a sequence of measures μ_n , ($n = 1, 2, 3, \dots$) with compact support lying in $F \cap U$, and having $\|\mu_n\| \leq \lambda(A)$ such that $\mu_n \rightarrow \lambda|A$, weak-star in $H^\infty(U)^*$.

In case U is the unit disk, this theorem is due to Stray [4]

Note that the equivalence of (1) and (2) does *not* say that “there is no interaction between distinct components in this problem”. It takes this neat form because of the formulation in terms of *closed* sets F .

We observe that (5) is equivalent to the same statement, with “closed set” replaced by “Borel set”.

We note that it is *not* true that (U, F) is Farrell if and only if (U^*, F) is Farrell. The latter condition implies the former, but not conversely. The classical *cornucopia* (Figure 1) provides an example.

figure 1 here : The cornucopia.

An example with smoothly-bounded components is provided by the *budding disc* (Figure 2).

figure 2 here: The budding disc.

Letting U denote the union of the buds,

$$U = \bigcup_{p=2}^{\infty} \bigcup_{j=0}^{2^p} \{z \in \mathbf{C} : |z - a_{p,j}| < 2^{-2^p}\},$$

where

$$a_{p,j} = \left(1 + 2^{-2^p}\right) \exp(2\pi j / 2^p),$$

we find that U^* is obtained from U by adding the open unit disc. Taking

$$S = \{a_{p,j} : p \geq 2, 0 \leq j \leq 2^p\},$$

and $F = \text{clos}(S)$, we see that (U, F) is once again Farrell, while (U^*, F) is not.

2. Proof of Theorem 1.

We begin with a lemma about FRS sets.

Lemma 1. *If V is a union of components of an FRS set, then V is itself an FRS set.*

Proof. Let U be an FRS set, with components U_1, U_2, U_3, \dots , and let

$$V = \bigcup_{n \in J} U_n$$

be a union of components of U . We must prove that for each $n \in J$, the set U_n is a component of V^* .

Fix $n \in J$, and let W be the component of V^* that contains U_n .

If $W \neq U_n$, then there exists a point

$$a \in W \cap \text{bdy}(U_n) \subset V^* \cap \text{bdy } U_n.$$

But then $a \in \text{bdy}(U^*)$, since $\text{bdy}(U_n) \subset \text{bdy}(U^*)$ and U is FRS. Since $\mathbf{C} \sim \text{clos}(U^*)$ is a subset of $\mathbf{C} \sim V$, this implies that $a \in \text{clos}(\mathbf{C} \sim V^*)$, contradicting the openness of V^* .

Thus $W = U_n$. ■

Now we begin the proof of Theorem 1.

We use the notation $S = F \cap U$, $X_n = \text{bdy } U_n$, $F_n = F \cap \text{clos}(U_n)$, $S_n = F \cap U_n$.

(1) \Rightarrow (2): Suppose (U, F) is a Farrell pair. Fix n . Let $f \in H^\infty(U_n)$ and $\epsilon > 0$ be given. We have to show that we may approximate f pointwise on U_n by polynomials p_m having

$$\|p_m\|_{U_n} \leq \|f\|_{U_n}, \text{ and}$$

$$\|p_m\|_{F_n} \leq \|f\|_{S_n} + \epsilon.$$

Extend f to an element of $H^\infty(U)$ by defining $f = 0$ on the remaining components U_m ($m \neq n$) of U . Then $\|f\|_U = \|f\|_{U_n}$ and $\|f\|_F = \|f\|_{F_n}$, so the existence of the required p_m 's follows from the fact that (U, F) is Farrell.

(2) \Rightarrow (3): This implication may be proved component by component. Further, since, by Lemma 1, the components of an FRS set are themselves FRS sets, we may suppose that U is connected for this part of the proof.

So let U be a connected FRS set and $F \subset \text{clos}(U)$ be such that (U, F) is Farrell. We must show that the extension \tilde{f} in $L^\infty(\lambda)$ has $|\tilde{f}| \leq \|f\|_S$, λ -a.e.. This is implicitly proved in Stray's paper, in case $U = \mathbf{D}$, but we cannot simply reduce to that case, because the conformal map from U to \mathbf{D} may fail to extend continuously (FRS examples exist. See section 3, below.), hence there is no simple correspondence of Farrell pairs (U, F) and (\mathbf{D}, F') . Here is a direct argument, based on Stray's idea.

Fix $f \in H^\infty(U)$ and $\eta > \|f\|_S$.

Suppose (for a contradiction) that there exists a compact set $A \subset F$, with $\lambda(A) > 0$, on which $|f| > \eta$. By means of obvious modification to f , η and A , we may arrange that

$$\|f\|_S \leq \eta < 1 = \|f\|_U, \text{ and}$$

the real part $\Re f$ of f exceeds $\frac{\eta+1}{2}$ on A .

Let $p_m \in \mathbf{C}[z]$ be any sequence converging pointwise to f on U , with $\|p_m\|_U \leq 1$.

We claim that necessarily $\liminf \|p_m\|_F > \eta$.

By Fatou's Lemma, we may choose $z \in U$ such that the harmonic measure ν with respect to z of A exceeds $\frac{3+\eta}{4}$ and $\Re f(z) > \frac{\eta+1}{2}$.

Then, for large enough m , we get $|p_m(z)| > \frac{\eta+1}{2}$.

But

$$\begin{aligned} |p_m(z)| &\leq \int_A |p_m(\zeta)| d\nu(\zeta) + \int_{X \sim A} |p_m(\zeta)| d\nu(\zeta) \\ &\leq \|p_m\|_A + \frac{1-\eta}{4}, \\ \|p_m\|_A &> \frac{3\eta+1}{4}, \end{aligned}$$

and this yields the claim.

This contradicts the assumption that (U, F) is Farrell. Thus (2) implies (3).

(3) \Rightarrow (1): We resume the consideration of the general FRS set U . Suppose (3) holds. We must show that (U, F) is Farrell.

Let $f \in H^\infty(U)$ and $\epsilon > 0$ be given. Let $\eta = \|f\|_S + \epsilon$. We will modify an argument that goes back to Davie to show that f may be approximated pointwise on U by polynomials p_m that have $|p_m| \leq 1$ on U and $|p_m| \leq \eta$ on F .

First, we extend f to U^* by defining $f = 0$ on $U^* \sim U$. This makes f an element of $H^\infty(U^*)$.

Let λ^* denote a harmonic measure for U^* on X . Then $\lambda \ll \lambda^*$.

Let

$$N = \{g \in C(Y) : \|g\|_F \leq \eta, \|g\|_Y \leq 1\}.$$

Here $C(Y)$ denotes the uniform algebra of all continuous complex-valued functions on Y .

By Mergelyan's Theorem [2], the closure of $\mathbf{C}[z]$ in $C(Y)$ is the algebra $A(Y)$ of all those $g \in C(Y)$ that are analytic on U^* . Thus it is enough for us to show that f

belongs to the closure, in the topology of uniform convergence on compact subsets of U , of the set $N \cap A(Y)$. This is a locally-convex topology, so we may apply the Separation Theorem. Further, the continuous linear functionals on the space of analytic functions on U , with the topology of uniform convergence on compact sets, are represented by measures having compact support in U . Thus all we need show is that

$$\left| \int f d\mu \right| \leq 1$$

whenever μ is such a measure and satisfies

$$\left| \int h d\mu \right| \leq 1$$

for all $h \in N \cap A(Y)$.

Let μ have these properties. The set N is the unit ball for an equivalent norm on the space $C(Y)$. Thus we may apply the Hahn–Banach Theorem and the Riesz Representation Theorem to obtain a measure ν , supported on Y , such that

$$\begin{aligned} \left| \int g d\nu \right| &\leq 1, \quad \forall g \in N \\ \int h d\nu &= \int h d\mu, \quad \forall h \in N \cap A(Y). \end{aligned}$$

It follows that

$$|\nu|(Y \sim F) + \eta|\nu|(F) \leq 1.$$

We claim that $\nu|X$ is absolutely-continuous with respect to λ^* .

In fact, let μ_1 be the balayage of μ to X , and let ν_1 be the balayage of $\nu|U$. Then μ_1 and ν_1 are absolutely-continuous with respect to λ^* , and $\mu_1 - \nu_1 - (\nu|X)$ annihilates $A(Y)$. By a generalised F. and M. Riesz Theorem [2, Theorem (II.8.6)], $\mu_1 - \nu_1 - (\nu|X)$ is absolutely-continuous with respect to λ^* , hence so is $\nu|X$.

Thus \tilde{f} is defined ν -a.e., and we have that $|\tilde{f}| \leq \eta$ ν -a.e. on F , and $|\tilde{f}| \leq 1$, ν -a.e. on Y .

Choose $g_m \in A(Y)$ such that $g_m \rightarrow f$ pointwise λ^* -a.e. on $\text{bdy } Y$ and $\|g_m\|_Y \leq 1$. This is possible, by the Farrell–Rubel–Shields Theorem and the Reduction of Norm Theorem [2, (VI.5.3)].

Then

$$\begin{aligned} \left| \int f d\mu \right| &= \left| \int f d\nu \right| \\ &\leq \left| \int_{Y \sim F} f d\nu \right| + \left| \int_F f d\nu \right| \\ &\leq 1 \cdot |\nu|(Y \sim F) + \eta|\nu|(F) \leq 1, \end{aligned}$$

as required.

Thus (3) implies (1).

Before completing the proof, we give a second lemma.

Lemma 2. *Let $H \subset \mathbf{S}^1$ and $S \subset \mathbf{D}$ be Borel sets. Then the following are equivalent:*

- (1) *$d\theta$ -almost every point of H is a nontangential limit point of S .*
- (2) *For each Borel set $T \subset H$ there is a sequence of measures μ_n , ($n = 1, 2, 3, \dots$) with compact support lying in S , and having total mass less than or equal to the $d\theta$ -measure $|T|$ of T , such that $\mu_n \rightarrow d\theta|T$, weak-star in $H^\infty(\mathbf{D})^*$.*
- (3) *$|\tilde{f}| \leq \|f\|_S$ $d\theta$ -a.e. on H , for each $f \in H^\infty(\mathbf{D})$. Here \tilde{f} denotes the non-tangential limit of f .*

In formulating condition (2), we take the point of view that $d\theta|E$ acts on $H^\infty(\mathbf{D})$ via the usual (Fatou) map $f \mapsto \tilde{f}$, $H^\infty(\mathbf{D}) \rightarrow L^\infty(d\theta)$.

Note especially that condition (2) does *not* just say that $\mu_n \rightarrow d\theta|E$ in the usual weak-star topology on measures (i.e. weak-star in $C(\text{clos}(\mathbf{D}))$). That would not be sufficient, and (2) is a much stronger statement.

It is not hard to see that it is equivalent to demand condition (2) only for *closed* sets $T \subset H$.

Proof of Lemma.

(1) \Rightarrow (2): Assume that almost every point of H is a nontangential limit point of S , and let T be a closed subset of H .

For $\alpha > 0$, let us say that $z \in \mathbf{S}^1$ is an α -non-tangential limit point of S if there is a sequence $z_n \in S$, converging to z , such that

$$\alpha \cdot |z - z_n| \leq 1 - |z_n|.$$

We may assume that there exists $\alpha > 0$ such that each point of T is an α -non-tangential limit point of S , because the general T may be written as a disjoint union

$$T = \bigcup_{m=1}^{\infty} T_m,$$

where

$$\bigcup_{m=1}^p T_m$$

takes up all but the 2^{-p} -th part of the measure of T , and all points of T_m are α_m -non-tangential limit points of S for some $\alpha_m > 0$. Given suitable sequences $\nu_{m,n}$ for T_m , their term-by-term sum does the trick for T .

For $s \in S$, let W_s denote the arc of \mathbf{S}^1

$$W_s = \{z \in \mathbf{S}^1 : \alpha \cdot |s - z| \leq 1 - |s|\}.$$

For each n , take $r = (n - 1)/n$. Then $r < 1$, and the arcs W_s , corresponding to $s \in S$ with $|s| > r$, give a fine covering of T , so we may select points $s_m \in S$ ($m = 1, 2, 3, \dots$) with $|s_m| > r$ such that the corresponding W_{s_m} are pairwise disjoint and cover $d\theta$ -almost-all of T . Choose an integer M such that

$$\bigcup_{m=1}^M W_{s_m}$$

has all but a proportion $1 - r$ of the $d\theta$ -measure of T . Define ν_n to be the atomic measure having mass $|W_{s_m}|$ at the point s_m . Thus ν_n is a finite sum of point masses, supported on S .

It follows easily, using the fact that functions belonging to $H^\infty(\mathbf{D})$ have nontangential limits a.e. $d\theta$, that $\nu_n \rightarrow d\theta|T$ weak-star on $H^\infty(\mathbf{D})$.

(2) \Rightarrow (3): Suppose (2) holds. If some $f \in H^\infty(\mathbf{D})$ had $|\tilde{f}| > \|f\|_S$, then we could choose a closed set $T \subset H$, a unimodular constant ω and a positive ϵ such that $\Re(\omega f) > \|f\|_S + \epsilon$ on T . Taking ν_n as in (2), we would then have

$$\left| \int f d\nu_n - \int_T \tilde{f} d\theta \right| \geq \epsilon \cdot |T| > 0$$

for all n , contradicting the weak-star convergence.

(3) \Rightarrow (1): This was proved by Stray [4, p.302], in the course of proving his theorem. ■

We note from the proof that the measures ν_n may be taken discrete, if that is desired.

We now continue the proof of Theorem 1.

(4) is equivalent to (3). Since each function in $H^\infty(U_n)$ extends to a function in $H^\infty(U)$, condition (3) of the Theorem is equivalent to the condition that for each n and each $g \in H^\infty(\mathbf{D})$, the non-tangential boundary values \tilde{g} on \mathbf{S}^1 satisfy

$$|\tilde{g}| \leq \|g\|_{\phi_n(S_n)}, \quad d\theta\text{-a.e. on } H_n.$$

This is because with $f = g \circ \phi_n$, we get $\tilde{f} = \tilde{g} \circ \phi_n$ a.e. λ_n .

Thus the equivalence of conditions (3) and (4) of the Theorem follows from the equivalence of conditions (1) and (3) of Lemma 2.

(5) is equivalent to (3). In the same way, the equivalence of conditions (1) and (2) of Lemma 2 shows that condition (3) of the Theorem is equivalent to the statement that:

for each n and for each Borel set $T \subset H_n$ there is a sequence of measures μ_m , ($m = 1, 2, 3, \dots$) with compact support lying in $\phi_n(S_n)$, and having total mass less than or equal to the $d\theta$ -measure $|T|$ of T , such that $\mu_m \rightarrow d\theta|T$, weak-star in $H^\infty(\mathbf{D})^*$.

Since all the terms in this statement are conformally–invariant, we obtain the equivalence of condition (5) of the Theorem.

This concludes the proof of the Theorem.

3. Remarks and Further Results.

(3.1) Figure 3 shows an example of an FRS set U for which the conformal map from U to \mathbf{D} is not uniformly continuous.

figure 3 here.

(3.2) Let U be a FRS set and $F \subset \text{clos}(U)$ be closed.

From the proof of Theorem 1, we see that (U, F) is Farrell if and only if (U^*, F^*) is Farrell, where F^* is obtained from F by adding to it a set S_n in each component U_n of $U^* \sim U$ for which the harmonic measure w.r.t. U_n of F is positive, in such a way that

$\text{clos } S_n = S_n \cup (\text{bdy } U_n \cap F)$ and

$(U_n, \text{clos } S_n)$ is a Farrell set.

(3.3) Let $U = \bigcup U_n$ be an FRS set and $S \subset U$. If $(U, \text{clos } S)$ is Farrell, then there does not exist n such that

$$\lambda_n(\text{clos}(S) \sim \text{clos}(U_n \cap S)) > 0.$$

For if there were such an n , we could choose a function u , harmonic on U_n , with

$$u = 0 \text{ near } \text{clos}(U_n \cap S) \cap \text{bdy } (U_n),$$

$$u = 1 \text{ on a set of positive } \lambda_n \text{ measure on } \text{clos}(S), \text{ and}$$

$$0 < u < 1 \text{ on } U_n.$$

Defining $f = \exp(u + \iota u^*)$ on U_n , and $f = 0$ on the remaining components of U , we would have

$$\|f\|_U = 1, \|f\|_S = \|f\|_{S \cap U_n} < 1, \text{ but}$$

$|\tilde{f}| = 1$ on a set of positive λ measure. Thus condition (3) of Theorem 1 would fail, with $F = \text{clos}(S)$, hence $(U, \text{clos } S)$ is not Farrell.

(3.4) In practical terms, if it is desired to check that some pair is a Farrell pair, then (4) and (5) of Theorem 1 are the most readily verified.

It is not, in general, possible to phrase the conditions in terms of nontangential approach, in the original set U . Nontangential approach (approach from a sector) is not a conformally-invariant idea, and (as is known) there may be almost no points on $\text{bdy}(U_n)$ that are nontangentially approachable.

Condition (5) has the advantage that it does not refer to the conformal maps, about which little may be known. It has the disadvantage that weak-star convergence with respect to $H^\infty(U)$ may be difficult to check. In general, it is fair to say that the sets for which the conformal maps are ill-behaved are the same as those for which the weak-star convergence is hard to check, so there is little to choose between the two conditions.

(3.5) The property of being a Farrell pair is not topological, i.e. there exist pairs (U, F) and (V, G) , mapped one to the other by a global homeomorphism of \mathbf{C} , such that (U, F) is Farrell and (V, G) is not. To make such an example, modify the budding disc in such a way that the closure of the set of buds meets the unit circle (the boundary of the inner disc) only on a subset K that is perfect and has length zero. Take U to be the union of the central disc and the set of all the buds, and S to be the set of buds. Let $F = S \cup K$. Then (U, F) is a Farrell pair, since K has harmonic measure zero for all components of U . We may construct a homeomorphism of \mathbf{C} onto itself, of the form

$$\Psi : r \exp(\iota\theta) \mapsto r \exp(\iota\psi(\theta)),$$

where ψ is a periodic homeomorphism of \mathbf{R} onto itself, with the property that $\Psi(K)$ has positive length. The result is that $(\Psi(U), \Psi(K))$ is not Farrell, because condition (3) of Theorem 1 fails. In fact, one can see that the characteristic function of the set of buds cannot be suitably approximated by polynomials.

(3.6) If the boundary of U is locally-smooth off an exceptional set which for each n is the union of a set of Hausdorff dimension less than 1 and a countable union of

impressions of prime ends of U_n , then (by Makarov's Theorem [3]) each λ_n is mutually-absolutely continuous with respect to length (1-dimensional Hausdorff measure) and (by a well-known result on conformal mapping) nontangentiality transfers faithfully under the conformal map ϕ_n , so we can then rephrase the property of being a Farrell pair in terms of nontangential limits. This covers all but extremely pathological examples. However, it must be said that extremely pathological examples are generic in plane topology. For instance, one can find three simply-connected open sets with the *same* boundary.

(3.7) Another consequence of Makarov's Theorem is that if F is a compact set of Hausdorff dimension less than 1, then (U, K) is a Farrell pair whenever U is a FRS set whose closure contains K .

(3.8) The property of being a Farrell pair can be rephrased in terms of the maximal ideal space M (or character space) of the uniform algebra $H^\infty(U)$. Here is a brief summary of the relevant facts. There is a continuous projection $\pi : M \rightarrow \mathbf{C}$ defined by $\pi(\phi) = \phi(z \mapsto z)$. The algebra $H^\infty(U)$ is isometrically isomorphic to the l^∞ orthogonal direct sum of the uniform algebras $H^\infty(U_n)$. For each n , we may consider the maximal ideal space M_n of $H^\infty(U_n)$. There is usually more to M than the disjoint union of the M_n , but the rest of it will not concern us.

The harmonic measure λ_n (for the fixed point $a_n \in U_n$) on X_n lifts to M_n in many ways, but in one special way. The Shilov boundary Sh_n of M_n may be identified with the Choquet boundary of the space of bounded harmonic functions on U_n ; there is a unique probability measure λ_n^* on Sh_n such that

$$\int_{\text{Sh}_n} h(x) d\lambda_n^*(x) = h(a_n)$$

whenever h is a bounded harmonic function on U_n ; evidently $\pi_* \lambda_n^* = \lambda_n$.

Theorem 2. *Under the hypotheses of Theorem 1, the following are equivalent:*

- (1) *The pair (U, F) is a Farrell pair;*

(2) For each $f \in H^\infty(U)$ and each n , we have

$$|f(\phi)| \leq \|f\|_{F \cap U}$$

for λ_n^* -almost-all $\phi \in \pi^{-1}(F)$;

(3) For each n , λ_n^* -almost-all points of $\pi^{-1}(F)$ belong to the weak-star closure in $H^\infty(U)^*$ of $U \cap F$.

Apart from the considerably enhanced level of abstraction, the proof of this is not essentially different from that of the theorems about λ_n 's.

Acknowledgment: This research was partially supported by EOLAS grant SC/90/070 and DGYICT Proyecto PS89-0135

References

- [1] S. D. FISHER. *Function Theory on Planar Domains*. (John Wiley & Sons, New York, 1983).
- [2] T. W. GAMELIN. Uniform Algebras. (Prentice-Hall, Englewood Cliffs, N.J, 1969).
- [3] N. G. MAKAROV. On the distortion of the boundary sets under conformal mapping. Proc. Lond. Math. Soc. (3) **51** (1985), 369–84.
- [4] A. STRAY. Pointwise bounded approximation by functions satisfying a side condition. Pacific J. Math. **51** (1974), 301–5.

Draft 5, 23/10/91.